

Table 2.70A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Tobacco Product Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	157,518	161,028	78,162	78,627	65,489	66,476
GEOGRAPHIC DIVISION						
Northeast	30,074	30,569	14,111	14,641	11,640	12,026
New England	8,431	8,450	3,801	4,004	3,118	3,139
Middle Atlantic	21,644	22,119	10,311	10,637	8,522	8,887
Midwest	37,792	38,447	19,424	19,311	16,448	16,533
East North Central	26,321	26,873	13,520	13,834	11,392	11,811
West North Central	11,471	11,574	5,904	5,477	5,055	4,723
South	56,638	57,714	28,784	29,172	24,401	25,012
South Atlantic	29,317	29,852	14,981	14,687	12,495	12,610
East South Central	10,067	10,467	5,117	5,490	4,602	4,797
West South Central	17,255	17,394	8,686	8,995	7,304	7,604
West	33,014	34,297	15,843	15,502	13,000	12,905
Mountain	10,128	9,899	4,916	4,631	4,080	3,926
Pacific	22,885	24,398	10,927	10,871	8,921	8,980
COUNTY TYPE						
Large Metro	73,958	74,727	36,061	35,362	29,526	29,175
Small Metro	49,919	51,423	24,572	25,538	20,738	21,816
250K - 1 Mil. Pop.	37,250	38,121	18,241	18,841	15,477	16,116
<250K Pop.	12,668	13,302	6,331	6,697	5,262	5,700
Nonmetro	33,642	34,878	17,530	17,726	15,225	15,485
Urbanized	10,016	10,151	4,945	5,143	4,223	4,396
Less Urbanized	19,787	20,970	10,612	10,698	9,229	9,389
Completely Rural	3,840	3,757	1,973	1,885	1,773	1,700

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.70B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Tobacco Product Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	70.5	71.4	35.0	34.8	29.3	29.5
GEOGRAPHIC DIVISION						
Northeast	70.2	70.9	33.0	34.0	27.2	27.9
New England	75.2	74.6	33.9	35.3	27.8	27.7
Middle Atlantic	68.5	69.6	32.6	33.5	27.0	28.0
Midwest	72.9	73.7	37.4	37.0	31.7	31.7
East North Central	72.5	73.3	37.2	37.8	31.4	32.2
West North Central	73.7	74.5	37.9	35.3	32.5	30.4
South	71.3	72.2	36.2	36.5	30.7	31.3
South Atlantic	71.4	72.0	36.5	35.4	30.4	30.4
East South Central	72.6 ^a	76.1	36.9	39.9	33.2	34.9
West South Central	70.4	70.4	35.4	36.4	29.8	30.8
West	67.2	68.0	32.3	30.7	26.5	25.6
Mountain	70.8	69.5	34.4	32.5	28.5	27.6
Pacific	65.7	67.4	31.4	30.0	25.6	24.8
COUNTY TYPE						
Large Metro	67.8	68.7	33.1	32.5	27.1	26.8
Small Metro	72.7	73.3	35.8	36.4	30.2	31.1
250K - 1 Mil. Pop.	73.0	73.2	35.8	36.2	30.3	30.9
<250K Pop.	71.8	73.7	35.9	37.1	29.8	31.6
Nonmetro	73.9	74.7	38.5	37.9	33.4	33.2
Urbanized	75.4	75.3	37.2	38.2	31.8	32.6
Less Urbanized	73.3	73.9	39.3	37.7	34.2	33.1
Completely Rural	72.9	77.4	37.4	38.8	33.6	35.0

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.71A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Tobacco Product Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	8,872	8,741	5,673	5,540	3,641	3,561
GEOGRAPHIC DIVISION						
Northeast	1,534	1,507	1,022	1,024	647	668
New England	439	414	294	297	189	199
Middle Atlantic	1,095	1,093	728	727	458	469
Midwest	2,140	2,185	1,420	1,422	941	970
East North Central	1,475	1,501	978	962	655	660
West North Central	665	684	442	459	286	310
South	3,344	3,244	2,111	2,056	1,379	1,295
South Atlantic	1,615	1,543	994	938	646	588
East South Central	658	606	415	387	299	274
West South Central	1,071	1,095	702	731	434	433
West	1,854	1,806	1,121	1,039	674	629
Mountain	635	596	412	381	259	240
Pacific	1,219	1,210	709	658	415	389
COUNTY TYPE						
Large Metro	3,944 ^a	3,639	2,449	2,290	1,493	1,412
Small Metro	2,791	2,882	1,804	1,822	1,210	1,171
250K - 1 Mil. Pop.	2,037	2,132	1,311	1,353	863	847
<250K Pop.	754	750	493	469	347	324
Nonmetro	2,137	2,221	1,420	1,428	938	978
Urbanized	574	576	369	397	245	268
Less Urbanized	1,284	1,388	856	867	554	586
Completely Rural	279	257	196	164	139	125

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.71B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Tobacco Product Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	38.0	37.0	24.3	23.5	15.6	15.1
GEOGRAPHIC DIVISION						
Northeast	36.5	35.4	24.3	24.0	15.4	15.7
New England	39.7	36.5	26.5	26.2	17.0	17.6
Middle Atlantic	35.4	35.0	23.5	23.3	14.8	15.0
Midwest	39.2	38.7	26.0	25.2	17.2	17.2
East North Central	39.1	38.3	25.9	24.6	17.4	16.9
West North Central	39.4	39.5	26.2	26.6	17.0	17.9
South	40.2	39.0	25.4	24.7	16.6	15.6
South Atlantic	39.7	37.2	24.4	22.6	15.9	14.2
East South Central	46.2	43.7	29.1	27.9	21.0	19.7
West South Central	37.9	39.5	24.8	26.4	15.3	15.6
West	34.4	33.5	20.8	19.3	12.5	11.7
Mountain	39.1	36.5	25.4	23.3	16.0	14.7
Pacific	32.3	32.2	18.8	17.5	11.0	10.4
COUNTY TYPE						
Large Metro	34.7 ^a	32.4	21.5	20.4	13.1	12.6
Small Metro	38.7	39.2	25.0	24.8	16.8	15.9
250K - 1 Mil. Pop.	37.8	38.9	24.3	24.7	16.0	15.4
<250K Pop.	41.6	40.1	27.2	25.1	19.1	17.3
Nonmetro	44.5	44.3	29.6	28.5	19.5	19.5
Urbanized	43.3	41.6	27.8	28.6	18.5	19.3
Less Urbanized	44.6	44.6	29.7	27.9	19.3	18.8
Completely Rural	46.8	49.4	32.8	31.7	23.3	24.1

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.72A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Tobacco Product Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	20,910	21,692	15,117	15,647	12,424	12,951
GEOGRAPHIC DIVISION						
Northeast	3,695	3,773	2,730	2,802	2,260	2,318
New England	1,020	1,032	768	747	633	606
Middle Atlantic	2,675	2,741	1,962	2,055	1,627	1,712
Midwest	5,408	5,378	4,101	3,975	3,436	3,309
East North Central	3,708	3,692	2,812	2,719	2,388	2,286
West North Central	1,700	1,686	1,288	1,256	1,048	1,023
South	7,397	7,703	5,257	5,487	4,328	4,626
South Atlantic	3,521	3,655	2,546	2,633	2,095	2,220
East South Central	1,393	1,434	997	996	865	850
West South Central	2,483	2,614	1,714	1,858	1,368	1,556
West	4,410	4,838	3,029	3,383	2,400	2,699
Mountain	1,436	1,465	1,050	1,057	868	855
Pacific	2,974	3,374	1,980	2,326	1,532	1,844
COUNTY TYPE						
Large Metro	9,580	9,706	6,803	6,976	5,592	5,696
Small Metro	7,068	7,368	5,150	5,235	4,177	4,343
250K - 1 Mil. Pop.	4,867	5,124	3,516	3,673	2,875	3,059
<250K Pop.	2,200	2,244	1,634	1,562	1,302	1,284
Nonmetro	4,263 ^a	4,618	3,164 ^a	3,436	2,655 ^a	2,911
Urbanized	1,410	1,572	1,025	1,183	840	971
Less Urbanized	2,470	2,665	1,855	1,967	1,581	1,679
Completely Rural	382	380	284	286	234	261

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.72B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Tobacco Product Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	72.1 ^a	73.6	52.2	53.1	42.9	43.9
GEOGRAPHIC DIVISION						
Northeast	72.3 ^a	75.1	53.4	55.8	44.2	46.1
New England	77.4	79.8	58.3	57.7	48.0	46.8
Middle Atlantic	70.5	73.5	51.7	55.1	42.9	45.9
Midwest	78.3	78.0	59.4	57.7	49.8	48.0
East North Central	77.2	77.2	58.5	56.9	49.7	47.8
West North Central	80.9	79.8	61.3	59.4	49.9	48.4
South	71.9	73.5	51.1	52.3	42.1	44.1
South Atlantic	71.5	72.5	51.7	52.2	42.5	44.0
East South Central	76.1	77.9	54.4	54.1	47.2	46.2
West South Central	70.3	72.6	48.5	51.6	38.7 ^a	43.2
West	66.1	68.3	45.4	47.7	36.0	38.1
Mountain	71.3	71.2	52.1	51.4	43.1	41.6
Pacific	63.8	67.1	42.5	46.3	32.9	36.7
COUNTY TYPE						
Large Metro	68.9	70.8	49.0	50.9	40.2	41.6
Small Metro	74.5	75.0	54.3	53.3	44.0	44.2
250K - 1 Mil. Pop.	73.4	74.5	53.0	53.4	43.3	44.5
<250K Pop.	77.1	76.0	57.3 ^a	52.9	45.7	43.5
Nonmetro	76.1	77.5	56.5	57.7	47.4	48.9
Urbanized	75.0	76.6	54.5	57.6	44.7	47.3
Less Urbanized	76.3	77.7	57.3	57.3	48.9	48.9
Completely Rural	79.0	80.4	58.7	60.5	48.3	55.2

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.73A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Tobacco Product Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	127,735	130,594	57,372	57,440	49,424	49,964
GEOGRAPHIC DIVISION						
Northeast	24,845	25,290	10,360	10,815	8,732	9,040
New England	6,972	7,005	2,739	2,960	2,296	2,334
Middle Atlantic	17,874	18,284	7,621	7,855	6,437	6,706
Midwest	30,244	30,884	13,903	13,915	12,071	12,254
East North Central	21,138	21,680	9,730	10,153	8,350	8,865
West North Central	9,106	9,205	4,173	3,762	3,721	3,389
South	45,897	46,768	21,416	21,629	18,694	19,092
South Atlantic	24,181	24,655	11,441	11,116	9,754	9,802
East South Central	8,016	8,427	3,705	4,107	3,439	3,674
West South Central	13,701	13,686	6,270	6,406	5,502	5,616
West	26,749	27,653	11,693	11,081	9,926	9,578
Mountain	8,057	7,838	3,455	3,193	2,952	2,831
Pacific	18,692	19,815	8,238	7,887	6,974	6,747
COUNTY TYPE						
Large Metro	60,434	61,381	26,809	26,096	22,442	22,067
Small Metro	40,059	41,174	17,618	18,481	15,351	16,302
250K - 1 Mil. Pop.	30,346	30,866	13,414	13,815	11,739	12,211
<250K Pop.	9,714	10,308	4,204	4,667	3,612	4,091
Nonmetro	27,242	28,039	12,946	12,862	11,632	11,595
Urbanized	8,032	8,002	3,551	3,563	3,138	3,157
Less Urbanized	16,032	16,918	7,902	7,864	7,094	7,125
Completely Rural	3,178	3,120	1,493	1,434	1,400	1,313

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.73B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Tobacco Product Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	74.7	75.7	33.6	33.3	28.9	29.0
GEOGRAPHIC DIVISION						
Northeast	74.1	74.8	30.9	32.0	26.1	26.7
New England	79.4	78.6	31.2	33.2	26.1	26.2
Middle Atlantic	72.3	73.4	30.8	31.5	26.0	26.9
Midwest	76.6	77.9	35.2	35.1	30.6	30.9
East North Central	76.2	77.6	35.1	36.3	30.1	31.7
West North Central	77.3	78.8	35.4	32.2	31.6	29.0
South	75.4	76.5	35.2	35.4	30.7	31.2
South Atlantic	75.4	76.4	35.7	34.4	30.4	30.4
East South Central	75.5 ^a	80.1	34.9 ^a	39.0	32.4	34.9
West South Central	75.4	74.6	34.5	34.9	30.3	30.6
West	72.2	72.9	31.6	29.2	26.8	25.2
Mountain	75.5	74.3	32.4	30.3	27.7	26.8
Pacific	70.9	72.3	31.2	28.8	26.4	24.6
COUNTY TYPE						
Large Metro	72.1	73.2	32.0	31.1	26.8	26.3
Small Metro	77.1	77.8	33.9	34.9	29.5	30.8
250K - 1 Mil. Pop.	77.9	77.7	34.4	34.8	30.1	30.7
<250K Pop.	74.8	77.9	32.4	35.3	27.8	30.9
Nonmetro	77.5	78.5	36.8	36.0	33.1	32.4
Urbanized	79.7	79.7	35.2	35.5	31.2	31.5
Less Urbanized	76.8	77.5	37.8	36.0	34.0	32.6
Completely Rural	75.9	80.7	35.6	37.1	33.4	34.0

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.74A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Tobacco Product Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	148,646	152,286	72,489	73,087	61,848	62,915
GEOGRAPHIC DIVISION						
Northeast	28,540	29,062	13,090	13,617	10,993	11,358
New England	7,991	8,037	3,507	3,707	2,929	2,940
Middle Atlantic	20,549	21,025	9,583	9,910	8,064	8,418
Midwest	35,652	36,262	18,004	17,890	15,507	15,563
East North Central	24,846	25,372	12,542	12,872	10,737	11,151
West North Central	10,806	10,891	5,462	5,018	4,769	4,413
South	53,294	54,470	26,673	27,116	23,022	23,717
South Atlantic	27,702	28,310	13,987	13,749	11,849	12,022
East South Central	9,409	9,861	4,702	5,103	4,303	4,523
West South Central	16,183	16,300	7,984	8,264	6,870	7,172
West	31,159	32,491	14,722	14,464	12,326	12,277
Mountain	9,493	9,303	4,504	4,251	3,820	3,686
Pacific	21,666	23,188	10,218	10,213	8,506	8,591
COUNTY TYPE						
Large Metro	70,013	71,088	33,612	33,072	28,033	27,763
Small Metro	47,127	48,542	22,768	23,716	19,528	20,645
250K - 1 Mil. Pop.	35,213	35,990	16,930	17,488	14,613	15,270
<250K Pop.	11,914	12,552	5,838	6,228	4,915	5,376
Nonmetro	31,505	32,657	16,109	16,298	14,287	14,507
Urbanized	9,442	9,574	4,575	4,746	3,978	4,128
Less Urbanized	18,502	19,583	9,756	9,831	8,675	8,804
Completely Rural	3,561	3,500	1,777	1,721	1,634	1,575

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 2.74B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Tobacco Product Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 and 2001

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2000	2001	2000	2001	2000	2001
TOTAL	74.4 ^a	75.4	36.3	36.2	30.9	31.1
GEOGRAPHIC DIVISION						
Northeast	73.9	74.8	33.9	35.1	28.5	29.2
New England	79.1	78.8	34.7	36.3	29.0	28.8
Middle Atlantic	72.0	73.4	33.6	34.6	28.3	29.4
Midwest	76.8	77.9	38.8	38.4	33.4	33.4
East North Central	76.4	77.5	38.6	39.3	33.0	34.1
West North Central	77.9	78.9	39.4	36.4	34.4	32.0
South	74.9	76.0	37.5	37.9	32.4	33.1
South Atlantic	74.9	75.8	37.8	36.8	32.0	32.2
East South Central	75.6 ^b	79.8	37.8	41.3	34.6	36.6
West South Central	74.6	74.3	36.8	37.7	31.7	32.7
West	71.3	72.1	33.7	32.1	28.2	27.3
Mountain	74.8	73.8	35.5	33.7	30.1	29.2
Pacific	69.8	71.5	32.9	31.5	27.4	26.5
COUNTY TYPE						
Large Metro	71.6	72.9	34.4	33.9	28.7	28.5
Small Metro	76.7	77.3	37.1	37.8	31.8	32.9
250K - 1 Mil. Pop.	77.2	77.2	37.1	37.5	32.0	32.8
<250K Pop.	75.3	77.6	36.9	38.5	31.0	33.2
Nonmetro	77.3	78.3	39.5	39.1	35.1	34.8
Urbanized	79.0	79.2	38.3	39.3	33.3	34.1
Less Urbanized	76.7	77.5	40.4	38.9	36.0	34.8
Completely Rural	76.2	80.7	38.0	39.7	35.0	36.3

*Low precision; no estimate reported.

NOTE: Any tobacco product indicates use at least once of cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any tobacco use in the past year does not include use of pipe tobacco.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.